

SCHRÖDINGER TYPE OPERATORS ON GENERALIZED MORREY SPACES

PENGTAO LI, XIN WAN, AND CHUANGYUAN ZHANG

ABSTRACT. In this paper we introduce a class of generalized Morrey spaces associated with Schrödinger operator $L = -\Delta + V$. Via a pointwise estimate, we obtain the boundedness of the operators $V^{\beta_2}(-\Delta + V)^{-\beta_1}$ and their dual operators on these Morrey spaces.

CONTENTS

1. Introduction	1
2. Preliminaries	2
2.1. Schrödinger operator and the auxiliary function	2
2.2. Generalized Morrey spaces associated with L	3
2.3. Calderón-Zygmund operators	4
3. Riesz transforms and the commutators on $L_{\alpha,\theta,V}^{p,q,\lambda}(\mathbb{R}^n)$	4
4. Schrödinger type operators on $L_{\alpha,\theta,V}^{p,q,\lambda}(\mathbb{R}^n)$	9
5. Boundedness of the commutators on $L_{\alpha,\theta,V}^{p,q,\lambda}(\mathbb{R}^n)$	14
References	20

1. INTRODUCTION

The investigation of Schrödinger operators on the Euclidean space $\mathbb{R}^n$ with nonnegative potentials which belong to the reverse Hölder class has attracted attention of many authors. Shen [16] studied the Schrödinger operator $L = -\Delta + V$, assuming the nonnegative potential V belongs to the reverse Hölder class B_q , $q \geq \frac{n}{2}$. In [16], Shen proved the L^p -boundedness of the operators $(-\Delta + V)^{i\gamma}$, $\nabla^2(-\Delta + V)^{-1}$, $\nabla(-\Delta + V)^{-1/2}$ and $\nabla(-\Delta + V)^{-1}\nabla$. For further information, we refer the reader to Guo-Li-Peng [7], Liu [10], Liu-Huang-Dong [12], Liu-Wang-Dong [13], Tang-Dong [19], Yang-Yang-Zhou [20, 21] and the references therein.

The purpose of this paper is to generalize the results of Shen [16] and Sugano [18] to a class of Morrey spaces associated with L , denoted by $L_{\alpha,\theta,V}^{p,q,\lambda}(\mathbb{R}^n)$. See Definition 2.8 below. The significance of these spaces is that for particular choices of the parameters p , q , λ , θ and α , one obtains many classical function spaces. In particular,

$\theta = 0, \alpha = 0, p = q, 0 < \lambda < 1,$	Morrey space $L^{p,\lambda}(\mathbb{R}^n)$ [14]
$\theta = 0, p = q, 0 < \lambda < 1,$	Morrey type space $L_{\alpha,V}^{p,\lambda}(\mathbb{R}^n)$ [19]
$\alpha = \lambda = 0, \theta \in \mathbb{R}, 0 < p, q < \infty,$	Herz spaces $K_p^{\theta,q}$ [8]
$\alpha = 0, \lambda \geq 0, \theta \in \mathbb{R}, 0 < p, q < \infty,$	Morrey-Herz spaces $MK_{p,q}^{\theta,\lambda}$ [2, 15]

In Section 3, let T be one of the Schrodinger type operators $\nabla(-\Delta + V)^{-1}\nabla$, $\nabla(-\Delta + V)^{-1/2}$ and $(-\Delta + V)^{-1/2}\nabla$. With the help of the L^p -boundedness of T , it is easy to verify that T is bounded on $L_{\alpha,\theta,V}^{p,q,\lambda}(\mathbb{R}^n)$.

2010 *Mathematics Subject Classification.* Primary 42B35, 42B20.

Key words and phrases. Generalized Morrey spaces, Schrödinger operator, Commutator, reverse Hölder class.

For $b \in BMO(\mathbb{R}^n)$, we can also obtain the boundedness of the commutator $[b, T]$ on $L_{\alpha, \theta, V}^{p, q, \lambda}(\mathbb{R}^n)$. See Theorem 3.2 & 3.3. For $\theta = 0$, $p = q$ and $0 < \lambda < 1$, $L_{\alpha, 0, V}^{p, p, \lambda}(\mathbb{R}^n)$ becomes the spaces $L_{\alpha, V}^{p, \lambda}(\mathbb{R}^n)$ introduced by Tang-Dong [19]. Hence, the results are generalizations of [19, Theorems 1 & 2].

In recent years, the fractional integral operator $I_\alpha = (-\Delta + V)^{-\alpha}$ has been studied extensively. We refer to Duong-Yan [4], Jiang [6], Tang-Dong [19] and Yang-Yang-Zhou [20] for details. Suppose that $V \in B_s$, $s \geq \frac{n}{2}$. For $0 \leq \beta_2 \leq \beta_1 < \frac{n}{2}$, let

$$\begin{cases} T_{\beta_1, \beta_2} =: V^{\beta_2}(-\Delta + V)^{-\beta_1}, \\ T_{\beta_1, \beta_2}^* =: (-\Delta + V)^{-\beta_1} V^{\beta_2}. \end{cases}$$

Sugano [18] obtained the weighted estimates for $T_{\beta_1, \beta_2}, T_{\beta_1, \beta_2}^*$, $0 < \beta_2 \leq \beta_1 < 1$. If $\beta_2 = 0$, we can see that $T_{\beta_1, 0} = I_{\beta_1}$. So T_{β_1, β_2} and T_{β_1, β_2}^* can be seen as generalizations of I_α . Moreover, for $(\beta_1, \beta_2) = (1, 1)$ and $(1/2, 1/2)$, $T_{1, 1}^* = (-\Delta + V)^{-1}V$ and $T_{1/2, 1/2}^* = (-\Delta + V)^{-1/2}V^{1/2}$, respectively, which are studied by Shen [16] thoroughly. In Section 4, assume that $1 < p_1 < \infty$, $1 < p_2 < s/\beta_2$ and $1 < q < \infty$. If the index $(q, \beta_1, \beta_2, \lambda, \alpha, \theta)$ satisfies:

$$\begin{cases} 1/p_2 = 1/p_1 - 2(\beta_1 - \beta_2)/n, \\ \alpha \in (-\infty, 0] \text{ \& } \lambda \in (0, n), \\ \lambda/q - 1/p_1 + 2\beta_1/n < \theta < \lambda/q + 1 - 1/p_1, \end{cases}$$

we prove that T_{β_1, β_2} is bounded from $L_{\alpha, \theta, V}^{p_1, q, \lambda}(\mathbb{R}^n)$ to $L_{\alpha, \theta, V}^{p_2, q, \lambda}(\mathbb{R}^n)$. Specially, we know that $(-\Delta + V)^{-1}V$ and $(-\Delta + V)^{-1/2}V^{1/2}$ are bounded on $L_{\alpha, \theta, V}^{p, q, \lambda}(\mathbb{R}^n)$. See Theorems 4.7 & 4.8 for the details.

In the research of harmonic analysis and partial differential equations, the commutators play an important role. If T is a Calderón-Zygmund operator, $b \in BMO(\mathbb{R}^n)$, the L^p -boundedness of $[b, T]$ was first discovered by Coifman-Rochberg-Weiss [3]. Later, Strömberg [4] gave a simple proof, adopting the idea of relating commutators with the sharp maximal operator of Fefferman and Stein. In 2008, Guo-Li-Peng [7] introduced a condition $H(m)$ and obtained that L^p -boundedness of the commutator of Riesz transforms associated with L , where $b \in BMO(\mathbb{R}^n)$. For further information, we refer to Liu [11], Liu-Huang-Dong [12], Liu-Wang-Dong [13], Yang-Yang-Zhou [21] and the references therein.

In Section 5, by the boundedness of I_α and $(-\Delta + V)^{-\beta}V^{-\beta}$, we can deduce that the commutators $[b, T_{\beta_1, \beta_2}]$ and $[b, T_{\beta_1, \beta_2}^*]$ are bounded from $L^{p_1}(\mathbb{R}^n)$ to $L^{p_2}(\mathbb{R}^n)$. See Theorem 5.1. Theorem 5.1 together with Lemmas 4.1 & 2.7 can be used to prove that the commutators $[b, T_{\beta_1, \beta_2}]$ and $[b, T_{\beta_1, \beta_2}^*]$ are bounded from $L_{\alpha, \theta, V}^{p_1, q, \lambda}(\mathbb{R}^n)$ to $L_{\alpha, \theta, V}^{p_2, q, \lambda}(\mathbb{R}^n)$, respectively. See Theorems 5.2 & 5.3.

Remark 1.1. Unlike the setting of the Lebesgue spaces, it is well-known that the dual of $L^{p, \lambda}(\mathbb{R}^n)$ is not $L^{p', -\lambda}(\mathbb{R}^n)$. Hence, after obtaining Theorem 4.7, we can not deduce Theorem 4.8 via the method of duality used by Guo-Li-Peng [7].

2. PRELIMINARIES

2.1. Schrödinger operator and the auxiliary function. In this paper, we consider the Schrödinger differential operator $L = -\Delta + V$ on $\mathbb{R}^n$, $n \geq 3$, where V is a nonnegative potential belonging to the reverse Hölder class B_s , $s \geq \frac{n}{2}$, which is defined as follows.

Definition 2.1. Let V be a nonnegative function.

(i) We say $V \in B_s$, $s > 1$, if there exists $C > 0$ such that for every ball $B \subset \mathbb{R}^n$, the reverse Hölder inequality

$$\left(\frac{1}{|B|} \int_B V^s(x) dx \right)^{\frac{1}{s}} \lesssim \left(\frac{1}{|B|} \int_B V(x) dx \right)$$

holds.

(ii) We say $V \in B_\infty$ if there exists a constant C such that for every ball $B \subset \mathbb{R}^n$,

$$\|V\|_{L^\infty(B)} = \frac{1}{|B|} \int_B V(x) dx.$$

Remark 2.2. Assume $V \in B_s$, $1 < s < \infty$. Then $V(y)dy$ is a doubling measure. Namely, there exists a constant C_0 such that for any $r > 0$ and $y \in \mathbb{R}^n$,

$$(2.1) \quad \int_{B(x, 2r)} V(y) dy \lesssim C_0 \int_{B(x, r)} V(y) dy.$$

Definition 2.3. (Shen [16]) For $x \in \mathbb{R}^n$, the function $m_V(x)$ is defined as

$$\frac{1}{m_V(x)} =: \sup \left\{ r > 0 : \frac{1}{r^{n-2}} \int_{B(x, r)} V(y) dy \leq 1 \right\}.$$

Remark 2.4. The function m_V reflects the scale of V essentially, but behaves better. It is deeply studied in Shen [16] and play a crucial role in our proof. We list a property of m_V which will be used in the sequel and refer the reader to Guo-Li-Peng [7] for the details.

We state some notations and properties of m_V .

Lemma 2.5. ([16, Lemma 1.4]) Suppose that $V \in B_s$ with $s \geq \frac{n}{2}$. Then there exist positive constants C and k_0 such that

- (a) if $|x - y| \leq \frac{C}{m_V(x)}$, $m_V(x) \sim m_V(y)$;
- (b) $m_V(y) \lesssim (1 + |x - y|m_V(x))^{k_0} m_V(x)$;
- (c) $m_V(y) \geq C m_V(x) / \{1 + |x - y|m_V(x)\}^{k_0/(k_0+1)}$.

Lemma 2.6. ([16, Lemma 1.2]) Suppose that $V \in B_s$, $s > \frac{n}{2}$. There exists a constant C such that for $0 < r < R < \infty$,

$$\frac{1}{r^{n-2}} \int_{B(x, r)} V(y) dy \lesssim \left(\frac{R}{r}\right)^{\frac{n}{s}-2} \cdot \frac{1}{R^{n-2}} \int_{B(x, R)} V(y) dy.$$

Lemma 2.7. ([7, Lemma 2.3]) Suppose $V \in B_s$, $s > \frac{n}{2}$. Then for any $N > \log_2 C_0 + 1$, there exists a constant C_N such that for any $x \in \mathbb{R}^n$ and $r > 0$,

$$\frac{1}{(1 + r m_V(x))^N} \int_{B(x, r)} V(y) dy \lesssim C_N r^{n-2}.$$

2.2. Generalized Morrey spaces associated with L . Suppose that $V \in B_s$, $s > 1$. Let $L = -\Delta + V$ be the Schrödinger operator. Now we introduce a class of generalized Morrey spaces associated with L . For $k \in \mathbb{Z}$, let $E_k = B(x_0, 2^k r) \setminus B(x_0, 2^{k-1} r)$ and χ_k be the characteristic function of E_k .

Definition 2.8. Suppose that $V \in B_s$, $s > 1$. Let $p \in [1, +\infty)$, $q \in [1, +\infty)$, $\alpha \in (-\infty, +\infty)$ and $\lambda \in (0, n)$, $\theta \in (-\infty, +\infty)$. For $f \in L_{loc}^q(\mathbb{R}^n)$, we say $f \in L_{\alpha, \theta, V}^{p, q, \lambda}(\mathbb{R}^n)$ provided that

$$\|f\|_{L_{\alpha, \theta, V}^{p, q, \lambda}(\mathbb{R}^n)}^q = \sup_{B(x_0, r) \subset \mathbb{R}^n} \frac{(1 + r m_V(x_0))^\alpha}{r^{\lambda n}} \sum_{k=-\infty}^0 |E_k|^{\theta q} \|\chi_k f\|_{L^p(\mathbb{R}^n)}^q < \infty,$$

where $B(x_0, r)$ denotes a ball centered at x_0 and with radius r .

Proposition 2.9.

- (i) For $\alpha_1 > \alpha_2$, $L_{\alpha_1, \theta, V}^{p, q, \lambda}(\mathbb{R}^n) \subseteq L_{\alpha_2, \theta, V}^{p, q, \lambda}(\mathbb{R}^n)$;
- (ii) If $\theta = 0$, $p = q$ and $\alpha < 0$, $L^{p, \lambda}(\mathbb{R}^n) \subset L_{\alpha, \theta, V}^{p, q, \lambda}(\mathbb{R}^n)$;
- (iii) If $\theta = 0$, $p = q$ and $\alpha > 0$, $L_{\alpha, \theta, V}^{p, q, \lambda}(\mathbb{R}^n) \subset L^{p, \lambda}(\mathbb{R}^n)$.

2.3. Calderón-Zygmund operators. We say that an operator T taking $C_c^\infty(\mathbb{R}^n)$ into $L_{loc}^1(\mathbb{R}^n)$ is called a Calderón-Zygmund operator if

- (a) T extends to a bounded linear operator on $L^2(\mathbb{R}^n)$;
- (b) There exists a kernel K such that for every $f \in L_{loc}^1(\mathbb{R}^n)$;

$$Tf(x) = \int_{\mathbb{R}^n} K(x, y) f(y) dy \quad \text{a.e. on } \{\text{supp } f\}^c,$$

- (c) The kernel $K(x, y)$ satisfies the Calderón-Zygmund estimate

$$|K(x, y)| \leq \frac{C}{|x - y|^n};$$

$$|K(x + h, y) - K(x, y)| + |K(x, y + h) - K(x, y)| \leq \frac{C|h|^\delta}{|x - y|^{n+\delta}}$$

for $x, y \in \mathbb{R}^n$, $|h| < \frac{|x-y|}{2}$ and for some $\delta > 0$.

Shen [16] obtained the following result.

Theorem 2.10. [16, Theorem 0.8]) Suppose $V \in B_n$. Then

$$\nabla(-\Delta + V)^{-1}\nabla, \nabla(-\Delta + V)^{-\frac{1}{2}} \text{ and } (-\Delta + V)^{-\frac{1}{2}}\nabla$$

are Calderón-Zygmund operators.

Corollary 2.11. Suppose that $V \in B_n$ and $b \in BM(\mathbb{R}^n)$. The commutator $[b, T]$ is bounded on $L^p(\mathbb{R}^n)$.

In particular, let K denote the kernel of one of the above operators. Then K satisfies the following estimate:

$$(2.2) \quad |K(x, y)| \leq \frac{C_N}{(1 + |x - y| m_V(x))^N} \frac{1}{|x - y|^n}$$

for any $N \in \mathbb{N}$. See (6.5) of Shen [16] for the details.

Suppose $V \in B_s$ for $s \geq \frac{n}{2}$. Let $L = -\Delta + V$. The semigroup generated by L is defined as:

$$(2.3) \quad T_t f(x) = e^{-tL} f(x) = \int_{\mathbb{R}^n} K_t(x, y) f(y) dy, \quad f \in L^2(\mathbb{R}^n), \quad t > 0,$$

where K_t is the kernel of e^{-tL} .

Lemma 2.12. ([5]) Let $K_t(x, y)$ be as in (2.3). For every nonnegative integer k , there is a constant C_k such that

$$0 \leq K_t(x, y) \leq C_k t^{-\frac{n}{2}} \exp(-|x - y|^2/5t) (1 + \sqrt{t} m_V(x) + \sqrt{t} m_V(y))^{-k}.$$

Some notations. Throughout the paper, c and C will denote unspecified positive constants, possibly different at each occurrence. The constants are independence of the functions. $U \approx V$ represents that there is a constant $c > 0$ such that $c^{-1}V \leq U \leq cV$ whose right inequality is also written as $U \lesssim V$. Similarly, if $V \geq cU$, we denote $V \gtrsim U$.

3. RIESZ TRANSFORMS AND THE COMMUTATORS ON $L_{\alpha, \theta, V}^{p, q, \lambda}(\mathbb{R}^n)$

Throughout this paper, for $p \in (1, \infty)$, denote by p' the conjugate of p , that is, $\frac{1}{p} + \frac{1}{p'} = 1$. Let $V \in B_n$. In this section, we assume that T is one of the Schrodinger type operators $\nabla(-\Delta + V)^{-1}\nabla$, $\nabla(-\Delta + V)^{-1/2}$ and $(-\Delta + V)^{-1/2}\nabla$. We study the boundedness on $L_{\alpha, \theta, V}^{p, q, \lambda}(\mathbb{R}^n)$ of T and its commutator $[b, T]$ with $b \in BMO(\mathbb{R}^n)$. The bounded mean oscillation space $BMO(\mathbb{R}^n)$ is defined as follows.

Definition 3.1. A locally integrable function b is said to belong to $BMO(\mathbb{R}^n)$ if

$$\|b\|_{BMO} =: \sup_B \frac{1}{|B|} \int_B |b(x) - b_B| dx < \infty,$$

where the supremum is taken over all balls B in $\mathbb{R}^n$. Here $b_B = \frac{1}{|B|} \int_B b(x) dx$ stands for the mean value of b over the ball B and $|B|$ means the measure of B .

We first prove that T is bounded on $L_{\alpha, \theta, V}^{p, q, \lambda}(\mathbb{R}^n)$.

Theorem 3.2. Suppose that $\alpha \in (-\infty, 0]$, $\lambda \in (0, n)$ and $1 < q < \infty$. If $1 < p < \infty$, $\frac{\lambda}{q} - \frac{1}{p} < \theta < \frac{\lambda}{q} + 1 - \frac{1}{p}$, then the operators T are bounded on $L_{\alpha, \theta, V}^{p, q, \lambda}(\mathbb{R}^n)$.

Proof. For any ball $B(x_0, r)$, write

$$f(y) = \sum_{j=-\infty}^{\infty} f(y) \chi_j(y) = \sum_{j=-\infty}^{\infty} f_j(y),$$

where $E_j = B(x_0, 2^j r) \setminus B(x_0, 2^{j-1} r)$. Hence, we have

$$\begin{aligned} & (1 + rm_V(x_0))^\alpha r^{\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \|\chi_k T f\|_{L^p(\mathbb{R}^n)}^q \\ & \lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=-\infty}^{k-2} \|\chi_k T f_j\|_{L^p(\mathbb{R}^n)} \right)^q \\ & \quad + (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k-1}^{k+1} \|\chi_k T f_j\|_{L^p(\mathbb{R}^n)} \right)^q \\ & \quad + (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k+2}^{\infty} \|\chi_k T f_j\|_{L^p(\mathbb{R}^n)} \right)^q \\ & = A_1 + A_2 + A_3. \end{aligned}$$

For A_2 , by Theorem 2.10, we have

$$\begin{aligned} A_2 & \lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k-1}^{k+1} \|T f_j\|_{L^p(\mathbb{R}^n)} \right)^q \\ & \lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k-1}^{k+1} \|f_j\|_{L^p(\mathbb{R}^n)} \right)^q \\ & \lesssim \|f\|_{L_{\alpha, \theta, V}^{p, q, \lambda}}^q. \end{aligned}$$

We first estimate the term E_1 . Note that if $x \in E_k, y \in E_j$ and $j \leq k-2$, then $|x-y| \sim 2^k r$. By Lemma 2.5 and (2.2), we can get

$$\begin{aligned} \|\chi_k T f_j\|_{L^p(\mathbb{R}^n)} &\lesssim \left(\int_{E_k} \left| \int_{\mathbb{R}^n} \frac{1}{(1+|x-y| m_V(x))^N} \frac{1}{|x-y|^n} |f_j(y)| dy \right|^p dx \right)^{\frac{1}{p}} \\ &\lesssim \frac{1}{(1+2^k r m_V(x_0))^{N/k_0+1}} \frac{1}{(2^k r)^n} |E_k|^{\frac{1}{p}} \int_{E_j} |f(y)| dy \\ &\lesssim \frac{1}{(1+2^k r m_V(x_0))^{N/k_0+1}} |E_k|^{\frac{1}{p}-1} |E_j|^{\frac{1}{p'}} \left(\int_{E_j} |f(y)|^p dy \right)^{\frac{1}{p}}, \end{aligned}$$

where $\frac{1}{p} + \frac{1}{p'} = 1$. Since $-\frac{1}{p} + \frac{\lambda}{q} < \theta < (1 - \frac{1}{p}) + \frac{\lambda}{q}$, we obtain

$$\begin{aligned} A_1 &\lesssim (1 + r m_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=-\infty}^{k-2} \frac{|E_k|^{\frac{1}{p}-1} |E_j|^{\frac{1}{p'}} \|\chi_j f\|_{L^p(\mathbb{R}^n)}}{(1+2^k r m_V(x_0))^{N/k_0+1}} \right)^q \\ &\lesssim (1 + r m_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=-\infty}^{k-2} \frac{2^{\frac{n(j-k)}{p'}} (1+2^j r m_V(x_0))^{-\frac{\alpha}{q}}}{(1+2^k r m_V(x_0))^{N/k_0+1}} \right. \\ &\quad \times (2^j r)^{\frac{\lambda n}{q}} |E_j|^{-\theta} (1+2^j r m_V(x_0))^{\frac{\alpha}{q}} (2^j r)^{-\frac{\lambda n}{q}} (|E_j|^{\theta q} \|\chi_j f\|_{L^p(\mathbb{R}^n)}^q)^{\frac{1}{q}} \Big)^q \\ &\lesssim (1 + r m_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^\lambda \left(\sum_{j=-\infty}^{k-2} 2^{\frac{n(j-k)}{p'}} |E_k|^{\theta-\frac{\lambda}{q}} |E_j|^{\frac{\lambda}{q}-\theta} \right)^q \|f\|_{L_{\alpha,\theta,V}^{p,q,\lambda}(\mathbb{R}^n)}^q \\ &\lesssim (1 + r m_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^\lambda \left(\sum_{j=-\infty}^{k-2} 2^{(j-k)n(1-\frac{1}{p}+\frac{\lambda}{q}-\theta)} \right)^q \|f\|_{L_{\alpha,\theta,V}^{p,q,\lambda}(\mathbb{R}^n)}^q \\ &\lesssim \|f\|_{L_{\alpha,\theta,V}^{p,q,\lambda}(\mathbb{R}^n)}^q. \end{aligned}$$

For A_3 , we can see that when $x \in E_k, y \in E_j$, then $|x-y| \sim 2^j r$ for $j \geq k+2$. Similar to E_1 , we have

$$\begin{aligned} \|\chi_k T f_j\|_{L^p(\mathbb{R}^n)} &\lesssim \frac{1}{(1+2^j r m_V(x_0))^{N/k_0+1}} \frac{1}{(2^j r)^n} |E_k|^{\frac{1}{p}} \int_{E_j} |f(y)| dy \\ &\lesssim \frac{1}{(1+2^j r m_V(x_0))^{N/k_0+1}} \frac{1}{(2^j r)^n} |E_k|^{\frac{1}{p}} |E_j|^{\frac{1}{p'}} \left(\int_{E_j} |f(y)|^p dy \right)^{\frac{1}{p}} \\ &\lesssim \frac{1}{(1+2^j r m_V(x_0))^{N/k_0+1}} |E_k|^{\frac{1}{p}} |E_j|^{-\frac{1}{p}} \|\chi_j f\|_{L^p(\mathbb{R}^n)}. \end{aligned}$$

Since $-\frac{1}{p} + \frac{\lambda}{q} < \theta < (1 - \frac{1}{p}) + \frac{\lambda}{q}$, choosing N large enough, we obtain

$$\begin{aligned}
A_3 &\lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k+2}^{\infty} \frac{|E_k|^{\frac{1}{p}} |E_j|^{-\frac{1}{p}} \|\chi_j f\|_{L^p(\mathbb{R}^n)}}{(1 + 2^j rm_V(x_0))^{N/k_0+1}} \right)^q \\
&\lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left\{ \sum_{j=k+2}^{\infty} \frac{(1 + 2^j rm_V(x_0))^{-\frac{\alpha}{q}} (2^j r)^{\frac{\lambda n}{q}} |E_j|^{-\alpha}}{(1 + 2^j rm_V(x_0))^{N/k_0+1}} \right. \\
&\quad \times 2^{(k-j)\frac{n}{p}} (1 + 2^j rm_V(x_0))^{\frac{\alpha}{q}} (2^j r)^{-\frac{\lambda n}{q}} (|E_j|^{\theta q} \|\chi_j f\|_{L^p(\mathbb{R}^n)}^q)^{\frac{1}{q}} \Big\}^q \\
&\lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k+2}^{\infty} 2^{(k-j)\frac{n}{p}} |E_j|^{\frac{\lambda}{q}-\theta} \right)^q \|f\|_{L_{\alpha,\theta,V}^{p,q,\lambda}(\mathbb{R}^n)}^q \\
&\lesssim \|f\|_{L_{\alpha,\theta,V}^{p,q,\lambda}(\mathbb{R}^n)}^q.
\end{aligned}$$

Let $N = [-\frac{\alpha}{q} + 1](k_0 + 1)$. Finally, $\|Tf\|_{L_{\alpha,\theta,V}^{p,q,\lambda}(\mathbb{R}^n)} \lesssim \|f\|_{L_{\alpha,\theta,V}^{p,q,\lambda}(\mathbb{R}^n)}$. This completes the proof of Theorem 3.2. $\square$

Suppose that $b \in BMO(\mathbb{R}^n)$ and $V \in B_n$. Let T be one of the Schrodinger type operators $\nabla(-\Delta + V)^{-1}\nabla$, $\nabla(-\Delta + V)^{-1/2}$ and $(-\Delta + V)^{-1/2}\nabla$. The commutator $[b, T]$ is defined as

$$[b, T]f = bT(f) - T(bf).$$

Theorem 3.3. Suppose that $V \in B_n$ and $b \in BMO(\mathbb{R}^n)$. Let $1 < p < \infty$, $1 < q < \infty$, $\alpha \in (-\infty, 0]$, $\lambda \in (0, n)$. If the index (p, q, θ, λ) satisfies $\frac{\lambda}{q} - \frac{1}{p} < \theta < \frac{\lambda}{q} + 1 - \frac{1}{p}$, then

$$\|[b, T]f\|_{L_{\alpha,\theta,V}^{p,q,\lambda}} \leq C \|f\|_{L_{\alpha,\theta,V}^{p,q,\lambda}} \|b\|_{BMO}.$$

Proof. For any ball $B = B(x_0, r)$, we can get

$$f(y) = \sum_{j=-\infty}^{\infty} f(y) \chi_{E_j}(y) = \sum_{j=-\infty}^{\infty} f_j(y),$$

where $E_j = B(x_0, 2^j r) \setminus B(x_0, 2^{j-1} r)$. Hence, we have

$$\begin{aligned}
&(1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \|\chi_k [b, T]f\|_{L^p(\mathbb{R}^n)}^q \\
&\lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=-\infty}^{k-2} \|\chi_k [b, T]f_j\|_{L^p(\mathbb{R}^n)} \right)^q \\
&\quad + (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k-1}^{k+1} \|\chi_k [b, T]f_j\|_{L^p(\mathbb{R}^n)} \right)^q \\
&\quad + (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k+2}^{\infty} \|\chi_k [b, T]f_j\|_{L^p(\mathbb{R}^n)} \right)^q \\
&=: B_1 + B_2 + B_3.
\end{aligned}$$

For B_2 , by Corollary 2.11, we have

$$\begin{aligned}
B_2 &\lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k-1}^{k+1} \| [b, T] f_j \|_{L^p(\mathbb{R}^n)} \right)^q \\
&\lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k-1}^{k+1} \| f_j \|_{L^p(\mathbb{R}^n)} \right)^q \| b \|_{BMO}^q \\
&\lesssim \| f \|_{L_{\alpha, \theta, V}^{p, q, \lambda}}^q \| b \|_{BMO}^q.
\end{aligned}$$

Denote by $b_{2^k r}$ the mean value of b on the ball $B(x_0, 2^k r)$. For B_1 , by Lemma 2.5 and (2.2), we have

$$\begin{aligned}
&\| \chi_k [b, T] f_j \|_{L^p(\mathbb{R}^n)} \\
&\lesssim \frac{1}{(1 + 2^k rm_V(x_0))^{N/k_0+1}} \frac{1}{(2^k r)^n} \left[\int_{E_k} \left(\int_{E_j} |b(x) - b(y)| |f(y)| dy \right)^p dx \right]^{\frac{1}{p}} \\
&\lesssim \frac{1}{(1 + 2^k rm_V(x_0))^{N/k_0+1}} \frac{1}{(2^k r)^n} \left[\left(\int_{E_k} |b(x) - b_{2^k r}|^p dx \right)^{\frac{1}{p}} \int_{E_j} |f(y)| dy \right. \\
&\quad \left. + |E_k|^{\frac{1}{p}} \int_{E_j} |b(y) - b_{2^k r}| |f(y)| dy \right] \\
&\lesssim \frac{1}{(1 + 2^k rm_V(x_0))^{N/k_0+1}} \frac{1}{(2^k r)^n} \left[|E_k|^{\frac{1}{p}} |E_j|^{1-\frac{1}{p}} \| b \|_{BMO} \| f_j \|_{L^p(\mathbb{R}^n)} \right. \\
&\quad \left. + |E_k|^{\frac{1}{p}} \| f_j \|_{L^p(\mathbb{R}^n)} \left(\int_{E_j} |b(y) - b_{2^k r}|^{p'} dx \right)^{\frac{1}{p'}} \right] \\
&\lesssim \frac{1}{(1 + 2^k rm_V(x_0))^{N/k_0+1}} \frac{|E_j|^{1-\frac{1}{p}}}{|E_k|^{1-\frac{1}{p}}} (k-j) \| f_j \|_{L^p(\mathbb{R}^n)} \| b \|_{BMO},
\end{aligned}$$

where in the third inequality, we have used John-Nirenberg's inequality ([9]). Since $-\frac{1}{p} + \frac{\lambda}{q} < \theta < (1 - \frac{1}{p}) + \frac{\lambda}{q}$, we obtain

$$\begin{aligned}
B_1 &\lesssim \frac{(1 + rm_V(x_0))^\alpha}{r^{\lambda n}} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=-\infty}^{k-2} \frac{(k-j) \| f_j \|_{L^p(\mathbb{R}^n)}}{(1 + 2^k rm_V(x_0))^{N/k_0+1}} \frac{|E_j|^{1-\frac{1}{p}}}{|E_k|^{1-\frac{1}{p}}} \right)^q \| b \|_{BMO}^q \\
&\lesssim \frac{(1 + rm_V(x_0))^\alpha}{r^{\lambda n}} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left[\sum_{j=-\infty}^{k-2} \frac{(1 + 2^j rm_V(x_0))^{-\frac{\alpha}{q}}}{(1 + 2^k rm_V(x_0))^{N/k_0+1}} \| b \|_{BMO}^q \right. \\
&\quad \left. \times (k-j)(2^j r)^{\frac{\lambda n}{q}} |E_j|^{-\theta} \frac{|E_j|^{1-\frac{1}{p}}}{|E_k|^{1-\frac{1}{p}}} \right]^q \| f \|_{L_{\alpha, \theta, V}^{p, q, \lambda}}^q \\
&\lesssim \frac{(1 + rm_V(x_0))^\alpha}{r^{\lambda n}} \sum_{k=-\infty}^0 |E_k|^{\lambda} \left(\sum_{j=-\infty}^{k-2} (k-j) 2^{(k-j)n(\theta - \frac{\alpha}{q} + \frac{1}{p} - 1)} \right)^q \| f \|_{L_{\alpha, \theta, V}^{p, q, \lambda}}^q \| b \|_{BMO}^q \\
&\lesssim \| f \|_{L_{\alpha, \theta, V}^{p, q, \lambda}}^q \| b \|_{BMO}^q.
\end{aligned}$$

For B_3 , similar to B_1 , we have

$$\begin{aligned} & \|\chi_k[b, T]f_j\|_{L^p(\mathbb{R}^n)} \\ & \lesssim \frac{1}{(1 + 2^j r m_V(x_0))^{N/k_0+1}} \frac{1}{(2^j r)^n} \left(\int_{E_k} \left| \int_{E_j} |(b(x) - b(y))f(y)| dy \right|^p dx \right)^{\frac{1}{p}} \\ & \lesssim \frac{j - k}{(1 + 2^j r m_V(x_0))^{N/k_0+1}} |E_k|^{\frac{1}{p}} |E_j|^{-\frac{1}{p}} \|f_j\|_{L^p(\mathbb{R}^n)} \|b\|_{BMO}. \end{aligned}$$

Since $-\frac{1}{p} + \frac{1}{q} < \theta < (1 - \frac{1}{p}) + \frac{1}{q}$, choosing N large enough, we obtain

$$\begin{aligned} B_3 & \lesssim (1 + r m_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k+2}^{\infty} \frac{|E_k|^{\frac{1}{p}} |E_j|^{-\frac{1}{p}} (j - k) \|f_j\|_{L^p(\mathbb{R}^n)}}{(1 + 2^j r m_V(x_0))^{N/k_0+1}} \right)^q \|b\|_{BMO}^q \\ & \lesssim (1 + r m_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left[\sum_{j=k+2}^{\infty} \frac{(1 + 2^j r m_V(x_0))^{-\frac{\alpha}{q}}}{(1 + 2^j r m_V(x_0))^{N/k_0+1}} \right. \\ & \quad \left. \times (j - k) (2^j r)^{\frac{\lambda n}{q}} |E_j|^{-\theta} |E_k|^{\frac{1}{p}} |E_j|^{-\frac{1}{p}} \right]^q \|f\|_{L_{\alpha, \theta, V}^{p, q, \lambda}(\mathbb{R}^n)}^q \|b\|_{BMO}^q \\ & \lesssim (1 + r m_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^\lambda \left(\sum_{j=k+2}^{\infty} 2^{(k-j)n(\frac{1}{p} - \frac{1}{q} + \theta)} \right)^q \|f\|_{L_{\alpha, \theta, V}^{p, q, \lambda}(\mathbb{R}^n)}^q \|b\|_{BMO}^q \\ & \lesssim \|f\|_{L_{\alpha, \theta, V}^{p, q, \lambda}(\mathbb{R}^n)}^q \|b\|_{BMO}^q. \end{aligned}$$

Let $N = [-\frac{\alpha}{q} + 1](k_0 + 1)$. We finally get

$$\|[b, T]f\|_{L_{\alpha, \theta, V}^{p, q, \lambda}(\mathbb{R}^n)} \lesssim \|f\|_{L_{\alpha, \theta, V}^{p, q, \lambda}(\mathbb{R}^n)} \|b\|_{BMO}.$$

□

4. SCHRÖDINGER TYPE OPERATORS ON $L_{\alpha, \theta, V}^{p, q, \lambda}(\mathbb{R}^n)$

Let $L = -\Delta + V$ be the Schrödinger operator, where $V \in B_s$, $s > n/2$. For $0 < \beta < \frac{n}{2}$, the fractional integral operator associated with L is defined by

$$L^{-\beta}(f)(x) = \int_0^\infty e^{-tL}(f)(x) t^{\beta-1} dt.$$

Denote by $K_\beta(x, y)$ the kernel of $L^{-\beta}$. By Lemma 2.12, Bui [1] obtained the following pointwise estimate.

Lemma 4.1. ([1, Proposition 3.3]) *Let $0 < \beta < \frac{n}{2}$. For $N \in \mathbb{N}$, there is a constant C_N such that*

$$(4.1) \quad K_\beta(x, y) = \int_0^\infty K_t(x, y) t^{\beta-1} dt \leq \frac{C_N}{(1 + |x - y| m_V(x))^N} \frac{1}{|x - y|^{n-2\beta}},$$

where $K_t(\cdot, \cdot)$ is the kernel of the semigroup e^{-tL} .

Definition 4.2. Let $f \in L_{loc}^q(\mathbb{R}^n)$. Denote by $|B|$ the Lebesgue measure of the ball $B \subset \mathbb{R}^n$. The fractional Hardy-Littlewood maximal function $M_{\sigma, \gamma}$ is defined by

$$M_{\sigma, \gamma} f(x) = \sup_{x \in B} \left(\frac{1}{|B|^{1-\frac{\sigma \gamma}{n}}} \int_B |f(y)|^\gamma dy \right)^{\frac{1}{\gamma}}.$$

Lemma 4.3. ([3]) *Suppose $1 < \gamma < p_1 < \frac{n}{\sigma}$ and $\frac{1}{p_2} = \frac{1}{p_1} - \frac{\sigma}{n}$. Then*

$$\|M_{\sigma, \gamma} f\|_{L^{p_2}(\mathbb{R}^n)} \lesssim \|f\|_{L^{p_1}(\mathbb{R}^n)}.$$

As a generalization of the fractional integral associated with L , the operators $V^{\beta_2}(-\Delta + V)^{-\beta_1}$, $0 \leq \beta_2 \leq \beta_1 \leq 1$, have been studied by Sugano [18] systematically. Applying the method of Sugano [18] together with Lemma 4.1, we can obtain the following result for $V^{\beta_2}(-\Delta + V)^{-\beta_1}$, $0 \leq \beta_2 \leq \beta_1 \leq n/2$. We omit the proof.

Theorem 4.4. *Suppose that $V \in B_\infty$. Let $1 < \beta_2 \leq \beta_1 < \frac{n}{2}$. Then*

$$|V^{\beta_2}(-\Delta + V)^{-\beta_1} f(x)| \lesssim M_{2(\beta_1 - \beta_2), 1} f(x).$$

In a similar way, by (4.1), we can get the following estimate for the operators $(-\Delta + V)^{-\beta_1} V^{\beta_2}$, $0 \leq \beta_2 \leq \beta_1 < \frac{n}{2}$.

Theorem 4.5. *Suppose that $V \in B_s$ for $s > \frac{n}{2}$. Let $0 \leq \beta_2 \leq \beta_1 < \frac{n}{2}$. Then*

$$|(-\Delta + V)^{-\beta_1} (V^{\beta_2} f)(x)| \lesssim M_{2(\beta_1 - \beta_2), (\frac{s}{\beta_2})'} f(x),$$

where $(\frac{s}{\beta_2})'$ is the conjugate of $(\frac{s}{\beta_2})$.

Proof. Let $r = 1/m_V(x)$. By Lemma 4.1 and Hölder's inequality, we have

$$\begin{aligned} & |(-\Delta + V)^{-\beta_1} V^{\beta_2} f(x)| \\ & \lesssim \sum_{k=-\infty}^{\infty} \int_{2^{k-1}r \leq |x-y| \leq 2^k r} \frac{1}{(1 + 2^k r m_V(x_0))^N} \frac{1}{(2^k r)^{n-2\beta_1}} V(y)^{\beta_2} |f(y)| dy \\ & \lesssim \sum_{k=-\infty}^{\infty} \frac{(2^k r)^{2\beta_2}}{(1 + 2^k)^N} \left(\frac{1}{(2^k r)^n} \int_{B(x, 2^k r)} V(y) dy \right)^{\beta_2} M_{2(\beta_1 - \beta_2), (\frac{s}{\beta_2})'}(f)(x). \end{aligned}$$

For $k \geq 1$, because $V(y)dy$ is a doubling measure, we have

$$\begin{aligned} \frac{(2^k r)^2}{(2^k r)^n} \int_{B(x, 2^k r)} V(y) dy & \lesssim C_0^k \cdot 2^{(2-n)k} \frac{r^2}{r^n} \int_{B(x, r)} V(y) dy \\ & \lesssim (2^k)^{k_0}, \end{aligned}$$

where $k_0 = 2 - n + \log_2 C_0$. For $k \leq 0$, Lemma 2.6 implies that

$$\begin{aligned} \frac{(2^k r)^2}{(2^k r)^n} \int_{B(x, 2^k r)} V(y) dy & \lesssim \left(\frac{r}{2^k r} \right)^{\frac{n}{s}-2} \frac{r^2}{r^n} \int_{B(x, r)} V(y) dy \\ & \lesssim (2^k)^{2-\frac{n}{s}}. \end{aligned}$$

Taking N large enough, we get

$$|(-\Delta + V)^{-\beta_1} V^{\beta_2} f(x)| \lesssim M_{2(\beta_1 - \beta_2), (\frac{s}{\beta_2})'} f(x).$$

□

By Theorem 4.5 and the duality, we can obtain

Corollary 4.6. *Suppose $V \in B_s$ for $s > \frac{n}{2}$.*

(1) *If $1 < (\frac{s}{\beta_2})' < p_1 < \frac{n}{2\beta_1 - 2\beta_2}$ and $\frac{1}{p_2} = \frac{1}{p_1} - \frac{2\beta_1 - 2\beta_2}{n}$, then*

$$\|(-\Delta + V)^{-\beta_1} V^{\beta_2} f\|_{L^{p_2}(\mathbb{R}^n)} \lesssim \|f\|_{L^{p_1}(\mathbb{R}^n)},$$

where $\frac{s}{\beta_2} + (\frac{s}{\beta_2})' = 1$.

(2) *If $1 < p_2 < \frac{s}{\beta_2}$ and $\frac{1}{p_2} = \frac{1}{p_1} - \frac{2\beta_1 - 2\beta_2}{n}$, then*

$$\|V^{\beta_2}(-\Delta + V)^{-\beta_1} f\|_{L^{p_2}(\mathbb{R}^n)} \lesssim \|f\|_{L^{p_1}(\mathbb{R}^n)}.$$

Theorem 4.7. Suppose that $V \in B_s$, $s \geq \frac{n}{2}$, $\alpha \in (-\infty, 0]$, $\lambda \in (0, n)$. Let $1 < q < \infty$, $1 < \beta_2 \leq \beta_1 < \frac{n}{2}$ and $1 < p_2 < \frac{s}{\beta_2}$ with $\frac{1}{p_1} - \frac{1}{p_2} = \frac{2\beta_1 - 2\beta_2}{n}$. If $\frac{\lambda}{q} - \frac{1}{p_1} + \frac{2\beta_1}{n} < \theta < \frac{\lambda}{q} + 1 - \frac{1}{p_1}$, then

$$\|V^{\beta_2}(-\Delta + V)^{-\beta_1} f\|_{L_{\alpha, \theta, V}^{p_2, q, \lambda}} \lesssim \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}.$$

Proof. For any ball $B(x_0, r)$, write

$$f(y) = \sum_{j=-\infty}^{\infty} f(y)\chi_{E_j}(y) = \sum_{j=-\infty}^{\infty} f_j(y),$$

where $E_j = B(x_0, 2^j r) \setminus B(x_0, 2^{j-1} r)$. Hence, we have

$$\begin{aligned} & (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \|\chi_k V^{\beta_2}(-\Delta + V)^{-\beta_1} f\|_{L^{p_2}(\mathbb{R}^n)}^q \\ & \lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=-\infty}^{k-2} \|\chi_k V^{\beta_2}(-\Delta + V)^{-\beta_1} f_j\|_{L^{p_2}(\mathbb{R}^n)} \right)^q \\ & + (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k-1}^{k+1} \|\chi_k V^{\beta_2}(-\Delta + V)^{-\beta_1} f_j\|_{L^{p_2}(\mathbb{R}^n)} \right)^q \\ & + (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k+2}^{\infty} \|\chi_k V^{\beta_2}(-\Delta + V)^{-\beta_1} f_j\|_{L^{p_2}(\mathbb{R}^n)} \right)^q \\ & = M_1 + M_2 + M_3. \end{aligned}$$

We first estimate M_2 . For $1 < p_2 < \frac{s}{\beta_2}$, by (2) of Corollary 4.6, we can get

$$M_2 \lesssim \frac{(1 + rm_V(x_0))^\alpha}{r^{\lambda n}} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k-1}^{k+1} \|f_j\|_{L^{p_1}(\mathbb{R}^n)} \right)^q \lesssim \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}^q.$$

Now we deal with the terms M_1 and M_3 . We choose N large enough such that

$$(N/k_0 + 1) - (\log_2 C_0 + 1)\beta_2 + \alpha/q > 0$$

and take a positive $N_1 < (N/k_0 + 1) - (\log_2 C_0 + 1)\beta_2$. For M_1 , note that if $x \in E_k, y \in E_j$ and $j \leq k - 2$, then $|x - y| \sim 2^k r$. By Lemmas 4.1 & 2.7, we use Hölder's inequality to obtain

$$\begin{aligned} & \|\chi_k V^{\beta_2}(-\Delta + V)^{-\beta_1} f_j\|_{L^{p_2}(\mathbb{R}^n)} \\ & \lesssim \left(\int_{E_k} |V^{\beta_2}(x)| \int_{E_j} \frac{1}{(1 + |x - y| m_V(x))^N} \frac{1}{|x - y|^{n-2\beta_1}} f(y) dy |^{p_2} dx \right)^{\frac{1}{p_2}} \\ & \lesssim \frac{1}{(1 + 2^k rm_V(x_0))^{N/k_0+1}} \frac{1}{(2^k r)^{n-2\beta_1}} \int_{E_j} |f(y)| dy \left(\int_{E_k} |V(x)|^{\beta_2 p_2} dx \right)^{\frac{1}{p_2}} \\ & \lesssim \frac{|E_j|^{1-\frac{1}{p_1}} |E_k|^{\frac{1}{p_2}}}{(1 + 2^k rm_V(x_0))^{N/k_0+1}} \frac{1}{(2^k r)^{n-2\beta_1}} \|f_j\|_{L^{p_1}(\mathbb{R}^n)} \left(\frac{1}{|E_k|} \int_{E_k} V(x)^s dx \right)^{\frac{\beta_2}{s}} \\ & \lesssim \frac{|E_j|^{1-\frac{1}{p_1}} |E_k|^{\frac{1}{p_2}}}{(1 + 2^k rm_V(x_0))^{N/k_0+1}} \frac{1}{(2^k r)^{n-2\beta_1}} \|f_j\|_{L^{p_1}(\mathbb{R}^n)} \left(\frac{1}{|B_k|} \int_{B_k} V(x) dx \right)^{\beta_2} \\ & \lesssim \frac{1}{(1 + 2^k rm_V(x_0))^{N_1}} \frac{1}{(2^k r)^{n-2\beta_1+2\beta_2}} |E_k|^{\frac{1}{p_2}} |E_j|^{1-\frac{1}{p_1}} \|f_j\|_{L^{p_1}(\mathbb{R}^n)}, \end{aligned}$$

where $\frac{1}{p_1} - \frac{1}{p_2} = \frac{2\beta_1 - 2\beta_2}{n}$. Since $\frac{\lambda}{q} - \frac{1}{p_1} + \frac{2\beta_1}{n} < \theta < \frac{\lambda}{q} + 1 - \frac{1}{p_1}$, we obtain

$$\begin{aligned}
M_1 &\lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \\
&\quad \times \left(\sum_{j=-\infty}^{k-2} \frac{1}{(1 + 2^k rm_V(x_0))^{N_1}} \frac{1}{(2^k r)^{n-2\beta_1+2\beta_2}} |E_k|^{\frac{1}{p_2}} |E_j|^{1-\frac{1}{p_1}} \|f_j\|_{L^{p_1}(\mathbb{R}^n)} \right)^q \\
&\lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \\
&\quad \times \left(\sum_{j=-\infty}^{k-2} \frac{(1 + 2^j rm_V(x_0))^{-\frac{\alpha}{q}} (2^j r)^{\frac{\lambda n}{q}} |E_j|^{-\theta}}{(1 + 2^k rm_V(x_0))^{N_1}} \frac{1}{(2^k r)^{n-2\beta_1+2\beta_2}} |E_k|^{\frac{1}{p_2}} |E_j|^{1-\frac{1}{p_1}} \right)^q \|f\|_{L_{\alpha, V, \theta}^{p_1, \lambda, q}}^q \\
&\lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^\lambda \left(\sum_{j=-\infty}^{k-2} 2^{(j-k)n(\frac{\lambda}{q} - \theta - \frac{1}{p_1} + 1)} \right)^q \|f\|_{L_{\alpha, V, \theta}^{p_1, \lambda, q}}^q \\
&\lesssim \|f\|_{L_{\alpha, V, \theta}^{p_1, \lambda, q}}^q.
\end{aligned}$$

For M_3 , note that when $x \in E_k, y \in E_j$ and $j \geq k + 2$, then $|x - y| \sim 2^j r$. Similar to E_1 , we have

$$\begin{aligned}
&\|\chi_k V^{\beta_2} (-\Delta + V)^{-\beta_1} f_j\|_{L^{p_2}(\mathbb{R}^n)} \\
&\lesssim \frac{1}{(1 + 2^j rm_V(x_0))^{N/k_0+1}} \frac{1}{(2^j r)^{n-2\beta_1}} \int_{E_j} |f(y)| dy \left(\int_{E_k} |V(x)|^{\beta_2 p_2} dx \right)^{\frac{1}{p_2}} \\
&\lesssim \frac{1}{(1 + 2^j rm_V(x_0))^{N_1}} |E_j|^{\frac{2\beta_1}{n} - \frac{1}{p_1}} |E_k|^{\frac{1}{p_2} - \frac{2\beta_2}{n}} \|f_j\|_{L^{p_1}(\mathbb{R}^n)},
\end{aligned}$$

where $\frac{1}{p_1} - \frac{1}{p_2} = \frac{2\beta_1 - 2\beta_2}{n}$. Since $\frac{\lambda}{q} - \frac{1}{p_1} + \frac{2\beta_1}{n} < \theta < \frac{\lambda}{q} + 1 - \frac{1}{p_1}$, we obtain

$$\begin{aligned}
M_3 &\lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \\
&\quad \times \left(\sum_{j=k+2}^{\infty} \frac{1}{(1 + 2^j rm_V(x_0))^{N_1}} |E_j|^{\frac{2\beta_1}{n} - \frac{1}{p_1}} |E_k|^{\frac{1}{p_2} - \frac{2\beta_2}{n}} \|f_j\|_{L^{p_1}(\mathbb{R}^n)} \right)^q \\
&\lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \\
&\quad \times \left(\sum_{j=k+2}^{\infty} \frac{(1 + 2^j rm_V(x_0))^{-\frac{\alpha}{q}} (2^j r)^{\frac{\lambda n}{q}} |E_j|^{-\theta}}{(1 + 2^k rm_V(x_0))^{N_1}} \frac{|E_k|^{\frac{1}{p_2} - \frac{2\beta_2}{n}}}{|E_j|^{\frac{2\beta_1}{n} - \frac{1}{p_1}}} \right)^q \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}^q \\
&\lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^\lambda \left(\sum_{j=k+2}^{\infty} 2^{(k-j)n(\theta - \frac{\lambda}{q} + \frac{1}{p_1} + \frac{2\beta_1}{n})} \right)^q \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}^q \\
&\lesssim \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}^q.
\end{aligned}$$

Choosing N large enough, we obtain

$$\|V^{\beta_2} (-\Delta + V)^{-\beta_1} f\|_{L_{\alpha, \theta, V}^{p_2, q, \lambda}} \lesssim \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}.$$

□

Theorem 4.8. Suppose that $V \in B_s$, $s \geq \frac{n}{2}$, $\alpha \in (-\infty, 0]$, $\lambda \in (0, n)$ and $1 < q < \infty$. Let $0 < \beta_2 \leq \beta_1 < \frac{n}{2}$, $\frac{s}{s-\beta_2} < p_1 < \frac{n}{2\beta_1-2\beta_2}$ with $\frac{1}{p_2} = \frac{1}{p_1} - \frac{2\beta_1-2\beta_2}{n}$. If $\frac{\lambda}{q} - \frac{1}{p_2} < \theta < \frac{\lambda}{q} - \frac{1}{p_2} + 1 - \frac{2\beta_1}{n}$, then

$$\|(-\Delta + V)^{-\beta_1} V^{\beta_2} f\|_{L_{\alpha, \theta, V}^{p_2, q, \lambda}} \lesssim \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}.$$

Proof. For any ball $B(x_0, r)$, let $E_j = B(x_0, 2^j r) \setminus B(x_0, 2^{j-1} r)$. We can decompose f as follows.

$$f(y) = \sum_{j=-\infty}^{\infty} f(y) \chi_{E_j}(y) = \sum_{j=-\infty}^{\infty} f_j(y).$$

Similar to the proof of Theorem 4.7, we have

$$\begin{aligned} & (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \|\chi_k (-\Delta + V)^{-\beta_1} V^{\beta_2} f\|_{L^{p_2}(\mathbb{R}^n)}^q \\ & \lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=-\infty}^{k-2} \|\chi_k (-\Delta + V)^{-\beta_1} V^{\beta_2} f_j\|_{L^{p_2}(\mathbb{R}^n)} \right)^q \\ & + C(1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k-1}^{k+1} \|\chi_k (-\Delta + V)^{-\beta_1} V^{\beta_2} f_j\|_{L^{p_2}(\mathbb{R}^n)} \right)^q \\ & + C(1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k+2}^{\infty} \|\chi_k (-\Delta + V)^{-\beta_1} V^{\beta_2} f_j\|_{L^{p_2}(\mathbb{R}^n)} \right)^q \\ & = L_1 + L_2 + L_3. \end{aligned}$$

For L_2 , because $1 < \frac{s}{s-\beta_2} < p_1 < \frac{n}{2\beta_1-2\beta_2}$, we use Corollary 4.6 to obtain

$$L_2 \lesssim \frac{(1 + rm_V(x_0))^\alpha}{r^{\lambda n}} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k-1}^{k+1} \|f_j\|_{L^{p_1}(\mathbb{R}^n)} \right)^q \lesssim \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}^q.$$

For L_1 , we can see that if $x \in E_k$ and $y \in E_j$, then $|x - y| \sim 2^k r$ for $j \leq k - 2$. By Hölder's inequality and the fact that $V \in B_s$, we deduce from Lemmas 4.1 & 2.7 that

$$\begin{aligned} & \|\chi_k (-\Delta + V)^{-\beta_1} V^{\beta_2} f_j\|_{L^{p_2}(\mathbb{R}^n)}^q \\ & \lesssim \frac{1}{(1 + 2^k rm_V(x_0))^{N/k_0+1}} \frac{|E_k|^{\frac{1}{p_2}}}{(2^k r)^{n-2\beta_1}} \int_{E_j} V(x)^{\beta_2} |f(y)| dy \\ & \lesssim \frac{1}{(1 + 2^k rm_V(x_0))^{N/k_0+1}} \frac{|E_k|^{\frac{1}{p_2}}}{(2^k r)^{n-2\beta_1}} |E_j|^{1-\frac{1}{p_1}} \left(\frac{1}{|B_j|} \int_{B_j} V(x) dx \right)^{\beta_2} \|f_j\|_{L^{p_1}(\mathbb{R}^n)} \\ & \lesssim \frac{1}{(1 + 2^k rm_V(x_0))^{N_2}} \frac{|E_k|^{\frac{1}{p_2}}}{(2^k r)^{n-2\beta_1}} |E_j|^{1-\frac{1}{p_1}} (2^j r)^{-2\beta_2} \|f_j\|_{L^{p_1}(\mathbb{R}^n)}, \end{aligned}$$

where $\frac{1}{p_2} = \frac{1}{p_1} - \frac{2\beta_1 - 2\beta_2}{n}$ and $N_2 < (N/k_0 + 1) - (\log_2 C_0 + 1)\beta_2$. Since $\frac{\lambda}{q} - \frac{1}{p_2} < \theta < \frac{\lambda}{q} - \frac{1}{p_2} + 1 - \frac{2\beta_1}{n}$, we obtain

$$\begin{aligned}
L_1 &\lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \\
&\quad \times \left(\sum_{j=-\infty}^{k-2} \frac{1}{(1 + 2^k rm_V(x_0))^{N_2}} \frac{|E_k|^{\frac{1}{p_2}}}{(2^k r)^{n-2\beta_1}} |E_j|^{1-\frac{1}{p_1}} (2^j r)^{-2\beta_2} \|f_j\|_{L^{p_1}(\mathbb{R}^n)} \right)^q \\
&\lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \\
&\quad \left(\sum_{j=-\infty}^{k-2} \frac{(1 + 2^j rm_V(x_0))^{-\frac{\alpha}{q}} (2^j r)^{\frac{\lambda n}{q}} |E_j|^{-\theta} |E_k|^{\frac{1}{p_2}} |E_j|^{1-\frac{1}{p_1}}}{(1 + 2^k rm_V(x_0))^{N_2} (2^k r)^{n-2\beta_1} (2^j r)^{2\beta_2}} \right)^q \|f\|_{L_{\alpha,V,\theta}^{p_1,\lambda,q}}^q \\
&\lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^\lambda \left(\sum_{j=-\infty}^{k-2} 2^{(k-j)n(\theta-\frac{\lambda}{q}+\frac{1}{p_2}-1+\frac{2\beta_1}{n})} \right)^q \|f\|_{L_{\alpha,\theta,V}^{p_1,\lambda,q}}^q \\
&\lesssim \|f\|_{L_{\alpha,V,\theta}^{p_1,q,\lambda}}^q.
\end{aligned}$$

For L_3 , note that when $x \in E_k, y \in E_j$ and $j \geq k+2$, then $|x-y| \sim 2^j r$. Similar to E_1 , we have

$$\begin{aligned}
&\|\chi_k(-\Delta + V)^{-\beta_1} V^{\beta_2} f_j\|_{L^{p_2}(\mathbb{R}^n)}^q \\
&\lesssim \frac{1}{(1 + 2^j rm_V(x_0))^{N/k_0+1}} \frac{|E_k|^{\frac{1}{p_2}}}{(2^j r)^{n-2\beta_1}} \int_{E_j} V(x)^{\beta_2} |f(y)| dy \\
&\lesssim \frac{1}{(1 + 2^j rm_V(x_0))^{N_2}} \frac{|E_k|^{\frac{1}{p_2}}}{(2^j r)^{n-2\beta_1}} |E_j|^{1-\frac{1}{p_1}} (2^j r)^{-2\beta_2} \|f_j\|_{L^{p_1}(\mathbb{R}^n)},
\end{aligned}$$

where $\frac{1}{p_2} = \frac{1}{p_1} - \frac{2\beta_1 - 2\beta_2}{n}$, and $N_2 < (N/k_0 + 1) - (\log_2 C_0 + 1)\beta_2$. Since $\frac{\lambda}{q} - \frac{1}{p_2} < \theta < \frac{\lambda}{q} - \frac{1}{p_2} + 1 - \frac{2\beta_1}{n}$, we obtain

$$\begin{aligned}
L_3 &\lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \\
&\quad \times \left(\sum_{j=k+2}^{\infty} \frac{1}{(1 + 2^j rm_V(x_0))^{N_2}} \frac{|E_k|^{\frac{1}{p_2}}}{(2^j r)^{n-2\beta_1}} |E_j|^{1-\frac{1}{p_1}} (2^j r)^{-2\beta_2} \|f_j\|_{L^{p_1}(\mathbb{R}^n)} \right)^q \\
&\lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^\lambda \left(\sum_{j=k+2}^{\infty} 2^{(k-j)n(\theta-\frac{\lambda}{q}+\frac{1}{p_2})} \right)^q \|f\|_{L_{\alpha,\theta,V}^{p_1,q,\lambda}}^q \\
&\lesssim \|f\|_{L_{\alpha,\theta,V}^{p_1,q,\lambda}}^q.
\end{aligned}$$

Let N large enough. We finally get $\|(-\Delta + V)^{-\beta_1} V^{\beta_2} f\|_{L_{\alpha,\theta,V}^{p_2,q,\lambda}} \lesssim \|f\|_{L_{\alpha,\theta,V}^{p_1,q,\lambda}}$. $\square$

5. BOUNDEDNESS OF THE COMMUTATORS ON $L_{\alpha,\theta,V}^{p,q,\lambda}(\mathbb{R}^n)$

In this section, let $b \in BMO(\mathbb{R}^n)$. We consider the boundedness of commutators $[b, (-\Delta + V)^{-\beta_1} V^{\beta_2}]$ and its duality on the generalized Morrey spaces $L_{\alpha,\theta,V}^{p,q,\lambda}(\mathbb{R}^n)$. For this purpose, we prove the commutator $[b, (-\Delta)^{-\beta_1} V^{\beta_2}]$ is bounded from $L^{p_1}(\mathbb{R}^n)$ to $L^{p_2}(\mathbb{R}^n)$. For sake of simplicity, we denote by $b_{2^k r}$ the mean value of b on the ball $B(x_0, 2^k r)$.

Theorem 5.1. Suppose that $V \in B_s$, $s \geq \frac{n}{2}$ and $b \in BMO(\mathbb{R}^n)$.

(i) If $0 < \beta_2 \leq \beta_1 < \frac{n}{2}$, $\frac{s}{s-\beta_2} < p_1 < \frac{n}{2\beta_1-2\beta_2}$, $\frac{1}{p_2} = \frac{1}{p_1} - \frac{2\beta_1-2\beta_2}{n}$, then

$$\left\| [b, (-\Delta + V)^{-\beta_1} V^{\beta_2}] f \right\|_{L^{p_2}(\mathbb{R}^n)} \lesssim \|f\|_{L^{p_1}(\mathbb{R}^n)} \|b\|_{BMO}.$$

(ii) If $1 < p_2 < \frac{s}{\beta_2}$ and $\frac{1}{p_2} = \frac{1}{p_1} - \frac{2\beta_1-2\beta_2}{n}$, then

$$\left\| [b, V^{\beta_2} (-\Delta + V)^{-\beta_1}] f \right\|_{L^{p_2}(\mathbb{R}^n)} \lesssim \|f\|_{L^{p_1}(\mathbb{R}^n)}.$$

Proof. We only prove (i). (ii) can be obtained by duality. Because $\beta_2 \leq \beta_1$, we can decompose the operator $(-\Delta + V)^{-\beta_1} V^{\beta_2}$ as

$$(-\Delta + V)^{-\beta_1} V^{\beta_2} = (-\Delta + V)^{\beta_2-\beta_1} (-\Delta + V)^{-\beta_2} V^{\beta_2}.$$

Denote by $L^{\beta_2-\beta_1}$ and T_{β_2} the operators $(-\Delta + V)^{\beta_2-\beta_1}$ and $(-\Delta + V)^{-\beta_2} V^{\beta_2}$, respectively. Then we can get

$$\begin{aligned} & [b, (-\Delta + V)^{-\beta_1} V^{\beta_2}] f(x) \\ &= [b, (-\Delta + V)^{\beta_2-\beta_1} (-\Delta + V)^{-\beta_2} V^{\beta_2}] f(x) \\ &= b L^{\beta_2-\beta_1} T_{\beta_2} f(x) - L^{\beta_2-\beta_1} T_{\beta_2} (b f)(x) \\ &= b L^{\beta_2-\beta_1} T_{\beta_2} f(x) - L^{\beta_2-\beta_1} (b T_{\beta_2} f(x)) + L^{\beta_2-\beta_1} (b T_{\beta_2} f(x)) - L^{\beta_2-\beta_1} T_{\beta_2} (b f)(x) \\ &= [b, L^{\beta_2-\beta_1}] T_{\beta_2} f(x) + L^{\beta_2-\beta_1} [b, T_{\beta_2}] f(x). \end{aligned}$$

By (1) of Corollary 4.6, we can get

$$\begin{aligned} \left\| [b, (-\Delta + V)^{-\beta_1} V^{\beta_2}] f \right\|_{L^{p_2}(\mathbb{R}^n)} &\lesssim \left\| [b, L^{\beta_2-\beta_1}] T_{\beta_2} f \right\|_{L^{p_2}(\mathbb{R}^n)} + \left\| L^{\beta_2-\beta_1} [b, T_{\beta_2}] f \right\|_{L^{p_2}(\mathbb{R}^n)} \\ &\lesssim \left\| T_{\beta_2} f \right\|_{L^{p_1}(\mathbb{R}^n)} + \left\| [b, T_{\beta_2}] f \right\|_{L^{p_1}(\mathbb{R}^n)} \\ &\lesssim \|f\|_{L^{p_1}(\mathbb{R}^n)}. \end{aligned}$$

This completes the proof. $\square$

In the rest of this section, we prove the boundedness of the commutators $[b, V^{\beta_2} (-\Delta + V)^{-\beta_1}]$ and $[b, (-\Delta + V)^{-\beta_1} V^{\beta_2}]$ on $L_{\alpha, \theta, V}^{p_2, q, \lambda}(\mathbb{R}^n)$, respectively.

Theorem 5.2. Suppose that $V \in B_s$, $s \geq \frac{n}{2}$, $\alpha \in (-\infty, 0]$, and $\lambda \in (0, n)$. Let $1 < q < \infty$, $1 < \beta_2 \leq \beta_1 < \frac{n}{2}$ and $1 < p_2 < \frac{s}{\beta_2}$ with $\frac{1}{p_1} - \frac{1}{p_2} = \frac{2\beta_1-2\beta_2}{n}$. If $\frac{\lambda}{q} - \frac{1}{p_1} + \frac{2\beta_1}{n} < \theta < \frac{\lambda}{q} + 1 - \frac{1}{p_1}$, then for $b \in BMO(\mathbb{R}^n)$,

$$\left\| [b, V^{\beta_2} (-\Delta + V)^{-\beta_1}] f \right\|_{L_{\alpha, \theta, V}^{p_2, q, \lambda}} \lesssim \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}} \|b\|_{BMO}.$$

Proof. For any ball $B(x_0, r)$, we have

$$f(y) = \sum_{j=-\infty}^{\infty} f(y) \chi_{E_j}(y) = \sum_{j=-\infty}^{\infty} f_j(y),$$

where $E_j = B(x_0, 2^j r) \setminus B(x_0, 2^{j-1} r)$. Hence, we have

$$\begin{aligned}
& (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left\| \chi_k [b, V^{\beta_2}(-\Delta + V)^{-\beta_1}] f \right\|_{L^{p_2}(\mathbb{R}^n)}^q \\
& \lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=-\infty}^{k-2} \left\| \chi_k [b, V^{\beta_2}(-\Delta + V)^{-\beta_1}] f_j \right\|_{L^{p_2}(\mathbb{R}^n)} \right)^q \\
& + (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k-1}^{k+1} \left\| \chi_k [b, V^{\beta_2}(-\Delta + V)^{-\beta_1}] f_j \right\|_{L^{p_2}(\mathbb{R}^n)} \right)^q \\
& + (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k+2}^{\infty} \left\| \chi_k [b, V^{\beta_2}(-\Delta + V)^{-\beta_1}] f_j \right\|_{L^{p_2}(\mathbb{R}^n)} \right)^q \\
& =: D_1 + D_2 + D_3.
\end{aligned}$$

For D_2 , by (ii) of Theorem 5.1, we have

$$\begin{aligned}
D_2 & \lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k-1}^{k+1} \|f_j\|_{L^{p_1}(\mathbb{R}^n)} \right)^q \|b\|_{BMO}^q \\
& \lesssim \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}^q \|b\|_{BMO}^q.
\end{aligned}$$

For D_1 , by Lemmas 2.7 & 4.1, we obtain

$$\begin{aligned}
& \left\| \chi_k [b, V^{\beta_2}(-\Delta + V)^{-\beta_1}] f_j \right\|_{L^{p_2}(\mathbb{R}^n)} \\
& \lesssim \frac{1}{(1 + 2^k rm_V(x_0))^{N/k_0+1}} \frac{1}{(2^k r)^{n-2\beta_1}} \left(\int_{E_k} \left| \int_{E_j} V^{\beta_2}(x) (b(x) - b(y)) f(y) dy \right|^{p_2} dx \right)^{\frac{1}{p_2}} \\
& \lesssim \frac{1}{(1 + 2^k rm_V(x_0))^{N/k_0+1}} \frac{1}{(2^k r)^{n-2\beta_1}} \left[\left(\int_{E_k} V^{\beta_2 p_2}(x) |b(x) - b_{2^k r}|^{p_2} dx \right)^{\frac{1}{p_2}} \int_{E_j} |f(y)| dy \right. \\
& \quad \left. + \left(\int_{E_k} V^{\beta_2 p_2}(x) dx \right)^{\frac{1}{p_2}} \int_{E_j} |b(y) - b_{2^k r}| |f(y)| dy \right] \\
& \lesssim \frac{\|b\|_{BMO}}{(1 + 2^k rm_V(x_0))^{N/k_0+1}} \frac{1}{(2^k r)^{n-2\beta_1}} \left[\left(\int_{E_k} V(x) dx \right)^{\beta_2} |E_k|^{\frac{1}{p_2} - \beta_2} \int_{E_j} |f(y)| dy \right. \\
& \quad \left. + \left(\int_{E_k} V(x) dx \right)^{\beta_2} |E_k|^{\frac{1}{p_2} - \beta_2} |E_j|^{1 - \frac{1}{p_1}} (k - j) \|f_j\|_{L^{p_1}(\mathbb{R}^n)} \right] \\
& \lesssim \frac{\|b\|_{BMO}}{(1 + 2^k rm_V(x_0))^{N_1}} \frac{k - j}{(2^k r)^{n-2\beta_1}} |E_k|^{\frac{1}{p_2} - \frac{2\beta_2}{n}} |E_j|^{1 - \frac{1}{p_1}} \|f_j\|_{L^{p_1}(\mathbb{R}^n)},
\end{aligned}$$

where $\frac{1}{p_1} - \frac{1}{p_2} = \frac{2\beta_1 - 2\beta_2}{n}$ and $N_1 < (N/k_0 + 1) - (\log_2 C_0 + 1)\beta_2$. Since $\frac{\lambda}{q} - \frac{1}{p_1} + \frac{2\beta_1}{n} < \theta < \frac{\lambda}{q} + 1 - \frac{1}{p_1}$, we obtain

$$\begin{aligned}
D_1 &\lesssim \|b\|_{BMO}^q (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \\
&\quad \times \left(\sum_{j=-\infty}^{k-2} \frac{1}{(1 + 2^k rm_V(x_0))^{N_1}} \frac{k-j}{(2^k r)^{n-2\beta_1}} |E_k|^{\frac{1}{p_2} - \frac{2\beta_2}{n}} |E_j|^{1 - \frac{1}{p_1}} \|f_j\|_{L^{p_1}(\mathbb{R}^n)} \right)^q \\
&\lesssim \|b\|_{BMO}^q (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \\
&\quad \times \left(\sum_{j=-\infty}^{k-2} \frac{(1 + 2^j rm_V(x_0))^{-\frac{\alpha}{q}} (2^j r)^{\frac{\lambda n}{q}} |E_j|^{-\theta} |E_k|^{\frac{1}{p_2} - \frac{2\beta_2}{n}}}{(1 + 2^k rm_V(x_0))^{N_1} (2^k r)^{n-2\beta_1} |E_j|^{\frac{1}{p_1} - 1}} (k-j) \right)^q \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}^q \\
&\lesssim \|b\|_{BMO}^q \frac{(1 + rm_V(x_0))^\alpha}{r^{\lambda n}} \sum_{k=-\infty}^0 |E_k|^\lambda \left(\sum_{j=-\infty}^{k-2} (k-j) 2^{(j-k)n(\frac{\lambda}{q} - \theta - \frac{1}{p_1} + 1)} \right)^q \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}^q \\
&\lesssim \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}^q \|b\|_{BMO}^q.
\end{aligned}$$

For D_3 , because $\frac{1}{p_1} - \frac{1}{p_2} = \frac{2\beta_1 - 2\beta_2}{n}$ and $N_1 < (N/k_0 + 1) - (\log_2 C_0 + 1)\beta_2$, we have

$$\begin{aligned}
&\|\chi_k[b, V^{\beta_2}(-\Delta + V)^{-\beta_1}] f_j\|_{L^{p_2}(\mathbb{R}^n)} \\
&\lesssim \frac{1}{(1 + 2^j rm_V(x_0))^{N/k_0 + 1}} \frac{1}{(2^j r)^{n-2\beta_1}} \left(\int_{E_k} \left| \int_{E_j} V(x)^{\beta_2} (b(x) - b(y)) f(y) dy \right|^{p_2} dx \right)^{\frac{1}{p_2}} \\
&\lesssim \frac{j-k}{(1 + 2^j rm_V(x_0))^{N_1}} |E_j|^{\frac{2\beta_1}{n} - \frac{1}{p_1}} |E_k|^{\frac{1}{p_2} - \frac{2\beta_2}{n}} \|b\|_{BMO} \|f_j\|_{L^{p_1}(\mathbb{R}^n)},
\end{aligned}$$

where we have used the fact that $|x - y| \sim 2^j r$ for $x \in E_k, y \in E_j$ and $j \geq k + 2$. Since $\frac{\lambda}{q} - \frac{1}{p_1} + \frac{2\beta_1}{n} < \theta < \frac{\lambda}{q} + 1 - \frac{1}{p_1}$, we obtain

$$\begin{aligned}
D_3 &\lesssim (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \\
&\quad \times \left(\sum_{j=k+2}^{\infty} \frac{j-k}{(1 + 2^j rm_V(x_0))^{N_1}} |E_j|^{\frac{2\beta_1}{n} - \frac{1}{p_1}} |E_k|^{\frac{1}{p_2} - \frac{2\beta_2}{n}} \|b\|_{BMO} \|f_j\|_{L^{p_1}(\mathbb{R}^n)} \right)^q \\
&\lesssim \|b\|_{BMO}^q (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \\
&\quad \times \left(\sum_{j=k+2}^{\infty} \frac{(1 + 2^j rm_V(x_0))^{-\frac{\alpha}{q}} (2^j r)^{\frac{\lambda n}{q}} |E_j|^{-\theta} |E_k|^{\frac{1}{p_2} - \frac{2\beta_2}{n}}}{(1 + 2^j rm_V(x_0))^{N_1} |E_j|^{\frac{2\beta_1}{n} - \frac{1}{p_1}}} (j-k) \right)^q \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}^q \\
&\lesssim \|b\|_{BMO}^q (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^\lambda \left(\sum_{j=k+2}^{\infty} (j-k) 2^{(k-j)n(\theta - \frac{\lambda}{q} + \frac{1}{p_1} + \frac{2\beta_1}{n})} \right)^q \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}^q \\
&\lesssim \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}^q \|b\|_{BMO}^q.
\end{aligned}$$

Let N large enough. Finally, we get

$$\left\| [b, V^{\beta_2}(-\Delta + V)^{-\beta_1}] f \right\|_{L_{\alpha, \theta, V}^{p_2, q, \lambda}} \lesssim \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}} \|b\|_{BMO}.$$

□

Theorem 5.3. Suppose that $V \in B_s$, $s \geq \frac{n}{2}$ and $b \in BMO(\mathbb{R}^n)$. Let $\alpha \in (-\infty, 0]$, $\lambda \in (0, n)$ and $1 < q < \infty$. If $0 < \beta_2 \leq \beta_1 < \frac{n}{2}$, $\frac{s}{s-\beta_2} < p_1 < \frac{n}{2\beta_1-2\beta_2}$, $\frac{1}{p_2} = \frac{1}{p_1} - \frac{2\beta_1-2\beta_2}{n}$, $\frac{\lambda}{q} - \frac{1}{p_2} < \theta < \frac{\lambda}{q} - \frac{1}{p_2} + 1 - \frac{2\beta_1}{n}$, then

$$\left\| [b, (-\Delta + V)^{-\beta_1} V^{\beta_2}] f \right\|_{L_{\alpha, \theta, V}^{p_2, q, \lambda}} \lesssim \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}} \|b\|_{BMO}.$$

Proof. Similarly, we can decompose f based on an arbitrary ball $B(x_0, r)$ as follows.

$$f(y) = \sum_{j=-\infty}^{\infty} f(y) \chi_{E_j}(y) = \sum_{j=-\infty}^{\infty} f_j(y),$$

where $E_j = B(x_0, 2^j r) \setminus B(x_0, 2^{j-1} r)$. Hence, we have

$$\begin{aligned} & (1 + rm_V(x_0))^{\alpha} r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left\| \chi_k [b, (-\Delta + V)^{-\beta_1} V^{\beta_2}] f \right\|_{L^{p_2}(\mathbb{R}^n)}^q \\ & \lesssim (1 + rm_V(x_0))^{\alpha} r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=-\infty}^{k-2} \left\| \chi_k [b, (-\Delta + V)^{-\beta_1} V^{\beta_2}] f_j \right\|_{L^{p_2}(\mathbb{R}^n)} \right)^q \\ & + (1 + rm_V(x_0))^{\alpha} r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k-1}^{k+1} \left\| \chi_k [b, (-\Delta + V)^{-\beta_1} V^{\beta_2}] f_j \right\|_{L^{p_2}(\mathbb{R}^n)} \right)^q \\ & + (1 + rm_V(x_0))^{\alpha} r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k+2}^{\infty} \left\| \chi_k [b, (-\Delta + V)^{-\beta_1} V^{\beta_2}] f_j \right\|_{L^{p_2}(\mathbb{R}^n)} \right)^q \\ & = F_1 + F_2 + F_3. \end{aligned}$$

Applying Theorem 5.1, we can get

$$\begin{aligned} F_2 & \lesssim \frac{(1 + rm_V(x_0))^{\alpha}}{r^{\lambda n}} \sum_{k=-\infty}^0 |E_k|^{\theta q} \left(\sum_{j=k-1}^{k+1} \|f_j\|_{L^{p_1}(\mathbb{R}^n)} \right)^q \|b\|_{BMO}^q \\ & \lesssim \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}^q \|b\|_{BMO}^q. \end{aligned}$$

For F_1 , by Hölder's inequality and the fact that $V \in B_s$, we apply Lemmas 4.1 & 2.7 to deduce that

$$\begin{aligned} & \left\| \chi_k [b, (-\Delta + V)^{-\beta_1} V^{\beta_2}] f_j \right\|_{L^{p_2}(\mathbb{R}^n)} \\ & \lesssim \frac{1}{(1 + 2^k rm_V(x_0))^{N/k_0+1}} \frac{1}{(2^k r)^{n-2\beta_1}} \left(\int_{E_k} \left| \int_{E_j} (b(x) - b(y)) V^{\beta_2}(y) f(y) dy \right|^{p_2} dx \right)^{\frac{1}{p_2}} \\ & \lesssim \frac{1}{(1 + 2^k rm_V(x_0))^{N/k_0+1}} \frac{1}{(2^k r)^{n-2\beta_1}} \left[\left(\int_{E_k} |b(x) - b_{2^k r}|^{p_2} dx \right)^{\frac{1}{p_2}} \int_{E_j} |V^{\beta_2}(y) f(y)| dy \right. \\ & \quad \left. + |E_k|^{\frac{1}{p_2}} \int_{E_j} |b(y) - b_{2^k r}| V^{\beta_2}(y) f(y) dy \right] \\ & \lesssim \frac{(\int_{E_j} V(y) dy)^{\beta_2}}{(1 + 2^k rm_V(x_0))^{N/k_0+1}} \frac{k-j}{(2^k r)^{n-2\beta_1}} |E_k|^{\frac{1}{p_2}} |E_j|^{1-\frac{1}{p_1}} \|b\|_{BMO} \|f_j\|_{L^{p_1}(\mathbb{R}^n)} \\ & \lesssim \|b\|_{BMO} \frac{k-j}{(1 + 2^k rm_V(x_0))^{N_2}} |E_k|^{\frac{1}{p_2} + \frac{2\beta_1}{n} - 1} |E_j|^{1-\frac{1}{p_1} - \frac{2\beta_2}{n}} \|f_j\|_{L^{p_1}(\mathbb{R}^n)}, \end{aligned}$$

where $\frac{1}{p_2} = \frac{1}{p_1} - \frac{2\beta_1 - 2\beta_2}{n}$ and $N_2 < (N/k_0 + 1) - (\log_2 C_0 + 1)\beta_2$. Since $\frac{\lambda}{q} - \frac{1}{p_2} < \theta < \frac{\lambda}{q} - \frac{1}{p_2} + 1 - \frac{2\beta_1}{n}$, we obtain

$$\begin{aligned}
F_1 &\lesssim \|b\|_{BMO}^q (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \\
&\quad \times \left(\sum_{j=-\infty}^{k-2} \frac{k-j}{(1 + 2^k rm_V(x_0))^{N_2}} |E_k|^{\frac{1}{p_2} + \frac{2\beta_1}{n} - 1} |E_j|^{1 - \frac{1}{p_1} - \frac{2\beta_2}{n}} \|f_j\|_{L^{p_1}(\mathbb{R}^n)} \right)^q \\
&\lesssim \|b\|_{BMO}^q (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \\
&\quad \left(\sum_{j=-\infty}^{k-2} \frac{(1 + 2^j rm_V(x_0))^{-\frac{\alpha}{q}}}{(1 + 2^k rm_V(x_0))^{N_2}} (2^j r)^{\frac{\lambda n}{q}} |E_j|^{-\theta} \frac{|E_j|^{1 - \frac{1}{p_2} - \frac{2\beta_1}{n}}}{|E_k|^{1 - \frac{1}{p_2} - \frac{2\beta_1}{n}}} (k-j) \right)^q \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}^q \\
&\lesssim \|b\|_{BMO}^q \frac{(1 + rm_V(x_0))^\alpha}{r^{\lambda n}} \sum_{k=-\infty}^0 |E_k|^\lambda \left(\sum_{j=-\infty}^{k-2} (k-j) 2^{(k-j)n(\theta - \frac{\lambda}{q} + \frac{1}{p_2} - 1 + \frac{2\beta_1}{n})} \right)^q \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}^q \\
&\lesssim \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}^q \|b\|_{BMO}^q.
\end{aligned}$$

For F_3 , note that when $x \in E_k, y \in E_j$ and $j \geq k + 2$, then $|x - y| \sim 2^j r$. Similar to F_1 , we have

$$\begin{aligned}
&\|\chi_k[b, (-\Delta + V)^{-\beta_1} V^{\beta_2}] f_j\|_{L^{p_2}(\mathbb{R}^n)} \\
&\lesssim \frac{1}{(1 + 2^j rm_V(x_0))^{N/k_0 + 1}} \frac{1}{(2^j r)^{n - 2\beta_1}} \left(\int_{E_k} \left| \int_{E_j} (b(x) - b(y)) V(y)^{\beta_2} f(y) dy \right|^{p_2} dx \right)^{\frac{1}{p_2}} \\
&\lesssim \frac{j - k}{(1 + 2^j rm_V(x_0))^{N_2}} |E_k|^{\frac{1}{p_2}} |E_j|^{-\frac{1}{p_2}} \|f_j\|_{L^{p_1}(\mathbb{R}^n)} \|b\|_{BMO},
\end{aligned}$$

where $\frac{1}{p_2} = \frac{1}{p_1} - \frac{2\beta_1 - 2\beta_2}{n}$ and $N_2 < (N/k_0 + 1) - (\log_2 C_0 + 1)\beta_2$. Since $\frac{\lambda}{q} - \frac{1}{p_2} < \theta < \frac{\lambda}{q} - \frac{1}{p_2} + 1 - \frac{2\beta_1}{n}$, we obtain

$$\begin{aligned}
F_3 &\lesssim \|b\|_{BMO}^q (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^{\theta q} \\
&\quad \times \left(\sum_{j=k+2}^{\infty} \frac{(1 + 2^j rm_V(x_0))^{-\frac{\alpha}{q}}}{(1 + 2^k rm_V(x_0))^{N_2}} (2^j r)^{\frac{\lambda n}{q}} |E_j|^{-\theta} \frac{|E_k|^{\frac{1}{p_2}}}{|E_j|^{\frac{1}{p_2}}} (j - k) \|f_j\|_{L^{p_1}(\mathbb{R}^n)} \right)^q \\
&\lesssim \|b\|_{BMO}^q (1 + rm_V(x_0))^\alpha r^{-\lambda n} \sum_{k=-\infty}^0 |E_k|^\lambda \left(\sum_{j=k+2}^{\infty} (j - k) 2^{(k-j)n(\theta - \frac{\lambda}{q} + \frac{1}{p_2})} \right)^q \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}^q \\
&\lesssim \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}}^q \|b\|_{BMO}^q.
\end{aligned}$$

Let N large enough. We finally get

$$\| [b, (-\Delta + V)^{-\beta_1} V^{\beta_2}] f \|_{L_{\alpha, \theta, V}^{p_2, q, \lambda}} \lesssim \|f\|_{L_{\alpha, \theta, V}^{p_1, q, \lambda}} \|b\|_{BMO}.$$

□

Competing interests: The authors declare that they have no competing interests.

Authors' contributions: All authors read and approved the final manuscript.

Acknowledgements: Project supported by NSFC No.11171203; New Teacher's Fund for Doctor Stations, Ministry of Education No.20114402120003; Guangdong Natural Science Foundation S2011040004131; Foundation for Distinguished Young Talents in Higher Education of Guangdong, China, LYM11063.

REFERENCES

- [1] Bui, T-A: Weighted estimates for commutators of some singular integrals related to Schrödinger operators. *Bulletin des Sciences Mathématiques*. **138**, 270-292 (2014)
- [2] Chen, X, Chen, J: Boundedness of sublinear operators on generalized Morrey spaces and its application. *Chinese Ann. Math. Ser. A*. **32**, 705-720 (2011)
- [3] Coifman, R, Rochberg, R, Weiss, G: Factorization theorems for Hardy space in several variable. *Ann. Math.* **103**, 611-635 (1988)
- [4] Duong, X, Yan, L: On Commutators of fractional integrals. *Proc. Amer. Math. Soc.* **132**, 3549-3557 (2004)
- [5] Dziubański, J, Zienkiewicz, J: H^p spaces for Schrödinger operator. *Fourier analysis and related topics. Banach Center Publ.* **56**, 45-53 (2002)
- [6] Jiang, Y: Endpoint estimates for fractional integral associated to Schrödinger operators on the Heisenberg groups. *Acta Math. Sci. B.* **31**, 993-1000 (2011)
- [7] Guo, Z, Li, P, Peng, L: L^p boundedness of commutators of Riesz Transforms associated to Schrödinger operator. *J. Math. Anal. Appl.* **341**, 421-432 (2008)
- [8] Herz, C: Lipschitz spaces and Bernstein's theorem on absolutely convergent Fourier transforms. *J. Math. Mech.* **18**, 283-323 (1968/1969)
- [9] John, F, Nirenberg, L: On function of bounded mean oscillation. *Comm. Pure Appl. Math.* **14**, 415-426 (1961)
- [10] Liu, Y: The weighted estimates for the operators $V^\alpha(-\Delta_G + V)^{-\beta}$ and $V^\alpha \nabla_G(-\Delta_G + V)^{-\beta}$ on the stratified Lie group G . *J. Math. Anal. Appl.* **349**, 235-244 (2009)
- [11] Liu, Y: Commutators of BMO functions and degenerate Schrödinger operators with certain nonnegative potentials. *Monatsh. Math.* **165**, 41-56 (2012)
- [12] Liu Y, Huang J, Dong J: Commutators of Calderón-Zygmund operators related to admissible functions on spaces of homogeneous type and applications to Schrödinger operators. *Sci. China Math.* 2013, **56**: 1895-1913.
- [13] Liu, Y, Wang, L, Dong, J: Commutators of higher order Riesz transform associated with Schrödinger operators. *J. Funct. Spaces Appl.* **2013**, Art. ID 842375, 15 pp (2013)
- [14] Peetre, J: On the theory of $\mathcal{L}_{p,\lambda}$ spaces. *J. Funct. Anal.* **4**, 71-87 (1969)
- [15] Lu, S, Xu, L: Boundedness of rough singular integral operators on the homogeneous Morrey-Herz spaces. *Hokkaido Math. J.* **34**, 299-314 (2005)
- [16] Shen, Z: L^p estimate for Schrödinger operator with certain potentials. *Ann. Inst. Fourier*. **45**, 513-546 (1995)
- [17] Stein, EM: *Harmonic Analysis: Real-variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton Univ. Press, Princeton, NJ, 1993.
- [18] Sugano, S: Estimates for the operators $V^\alpha(-\Delta + V)^{-\beta}$ and $V^\alpha \nabla(-\Delta + V)^{-\beta}$ with certain non-negative potentials V . *Tokyo J. Math.* **21**, 441-452 (1998)
- [19] Tang, L, Dong, J: Boundedness for some Schrödinger type operators on Morrey spaces related to certain nonnegative potentials. *J. Math. Anal. Appl.* **355**, 101-109 (2009)
- [20] Yang, DC, Yang, DY, Zhou, Y: Endpoint properties of localized Riesz transforms and fractional integrals associated to Schrödinger operators. *Potential Anal.* **30**, 271-300 (2009)
- [21] Yang, DC, Yang, DY, Zhou, Y: Localized BMO and BLO spaces on RD-spaces and applications to Schrödinger operators. *Commun. Pure Appl. Anal.* **9**, 779-812 (2010)

COLLEGE OF MATHEMATICS, QINGDAO UNIVERSITY, QINGDAO, SHANDONG 266071, CHINA

E-mail address: ptli@qdu.edu.cn

DEPARTMENT OF MATHEMATICS, SHANTOU UNIVERSITY, SHANTOU, GUANGDONG 515063, CHINA.

E-mail address: 13xwan@stu.edu.cn

DEPARTMENT OF MATHEMATICS, SHANTOU UNIVERSITY, SHANTOU, GUANGDONG 515063, CHINA.

E-mail address: 12cyzhang@stu.edu.cn